



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 9/00	A2	(11) International Publication Number: WO 00/06706	(43) International Publication Date: 10 February 2000 (10.02.00)
--	----	---	---

(21) International Application Number: PCT/EP99/05413	(22) International Filing Date: 28 July 1999 (28.07.99)	(30) Priority Data: 9816639.0 30 July 1998 (30.07.98) GB	(71) Applicant (for all designated States except AT/US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VER. WALTUNGSGESSELLSCHAFT MBH [AT/AT]; Brunnerstrasse 59, A-1230 Vienna (AT). (72) Inventors; and (75) Inventors/Applicants (for US only): REVENKOVA, Ekaterina [RU/CH]; Peter Rot-Strasse 76, CH-4058 Basel (CH). PASZOWSKI, Jerzy [PL/CH]; Blauenweg 10, CH-4224 Nenzlingen (CH). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).
<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, MY, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>			

(54) Title: MAP KINASE PHOSPHATASE MUTANT

(57) Abstract

The present invention relates to DNA encoding proteins contributing to the regulation of a plant's response to DNA damage. DNA according to the invention comprises an open reading frame encoding a protein characterized by a stretch of amino acids or component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO:3. Preferably the DNA encodes a MAP kinase phosphatase.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MR	Mauritania	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Cote d'Ivoire	KP	Democratic People's	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Sequence alignments of SEQ ID NO: 3 using commercially available computer programs such as BLASTP of the NCBI BLAST family of programs or TFASTA or Bestfit of the Wisconsin Package Software, all based on well known algorithms for sequence identity or similarity searches, reveal that stretches of SEQ ID NO: 3 (component sequences) having more than 100 and preferably between 120 to 250 amino acids length can show between

DNA according to the present invention comprises an open reading frame encoding a protein characterized by a stretch of amino acids or component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO: 3. The protein characterized by SEQ ID NO: 3 is tracked down with the help of a T-DNA tagged Arabidopsis mutant showing hypersensitivity to methyl methanesulfonate (MMS). Said hypersensitivity as well as an observed hypersensitivity to other DNA damaging treatments such as UV light is indicative of the proteins' involvement in the repair of DNA damage, or in signaling pathways implicated in the response to similar genotoxic stress. The mutant is also sensitive to elevated temperature and anti-oxidant N-acetylcysteine. The mutant is not sensitive to osmotic shock, increased salinity, oxidative stress or elevated ethylene levels. An important characteristic of the mutant is cell death in response to growth in small closed vessels. This phenotype can be complemented by addition of abscisic acid (ABA) to the growth media. Furthermore, the mutant is more sensitive to exogenously applied ABA compared with the wild type which supports the notion that the genes disclosed by the present invention (SEQ ID NO: 1) are involved in stress signaling mediated by ABA.

The present invention relates to DNA encoding proteins contributing to the regulation of a plant's response to abiotic stress and in particular genotoxic stress. Cells of all organisms have evolved a series of DNA repair pathways which counteract the deleterious effects of DNA damage and are triggered by intricate signal cascades. To be able to modify or improve DNA repair using gene technology it is necessary to identify key proteins involved in said pathways or cascades. Therefore it is the main object of the present invention to provide DNA comprising an open reading frame encoding such a key protein.

MAP kinase phosphatase mutant

20% and almost 40% sequence identity to aligned stretches of known phosphatases, particularly phosphotyrosine phosphatases, MAP kinase phosphatases or dual specificity phosphatases. Protein phosphatases are classified by their substrate specificities as either phosphoserine/threonine phosphatases (PSTPs) or phosphotyrosine phosphatases (PTPs). The dual specificity phosphatases (DSPs) dephosphorylate both phosphotyrosine and phosphoserine/threonine residues and represent a subfamily of PTPs. MAP kinase phosphatases (MKPs) belong to the family of DSPs. The sequence VHCCQGVSR (SEQ ID NO: 4) found in SEQ ID NO: 3 can be interpreted as corresponding to the mammalian sequence motif IHGXAGXXRS (SEQ ID NO: 5) defining the family of PTPs, wherein the Ile at the first position can be replaced by Val and the Ser at the last position can be replaced by Thr

The present invention defines a new protein family the members of which are characterized by component amino acid sequences of more than 100 amino acid length showing 40% or higher amino acid sequence identity to aligned component sequences of SEQ ID NO: 3. Preferably said component sequences are of more than 120, more than 160 or even more than 200 amino acids length. The amino acid sequence identity is preferably higher than 50% or even higher than 55%. Most preferred are identities higher than 70%.

An example of DNA according to the present invention is described in SEQ ID NO: 1. The amino acid sequence of the protein encoded is identical to SEQ ID NO: 3. After alignment a stretch of the protein having about 140 amino acids shows 36% sequence identity to the MKP-1 protein described by Sun et al (Cell 75: 487-493, 1993). The identity determined after alignment with MKP-2 and MKP-3 is determined as 34% and 26%, respectively. Thus, according to the present invention a protein family related to MAP kinase phosphatases can be defined the members of which after alignment of a stretch of more than 100 amino acids length show 40% or higher amino acid sequence identity to SEQ ID NO: 3. Preferably, the amino acid sequence identity is higher than 50% or even higher than 55%. When making multiple sequence alignments, certain algorithms can take into account sequence similarities, such as same net charge or comparable hydrophobicity/hydrophilicity of the individual amino acids, in addition to sequence identities. The resulting values of sequence similarities, as compared to sequence identities, can help to assign a protein to the correct protein family in border-line cases. Proteins of particular interest, within the scope of the present invention, are MAP kinase phosphatases the amino acid sequence of which comprises at least one of the following characteristic amino acid subsequences:

- (a) TSILYDVFDYFEDV (SEQ ID NO: 6)
- (b) FVHCQGVSRSST (SEQ ID NO: 7)
- (c) FVHC (SEQ ID NO: 8)
- (d) QGVSR (SEQ ID NO: 9)
- (e) YFKSD (SEQ ID NO: 10)

DNA encoding proteins belonging to the new protein family according to the present invention can be isolated from monocotyledonous and dicotyledonous plants. Preferred sources are corn, sugar beet, sunflower, winter oilseed rape, soybean, cotton, wheat, rice, potato, broccoli, cauliflower, cabbage, cucumber, sweet corn, daikon, garden beans, lettuce, melon, pepper, squash, tomato, or watermelon. However, they can also be isolated from mammalian sources such as mouse or human tissues. The following general method, can be used, which the person skilled in the art will normally adapt to his specific task. A single stranded fragment of SEQ ID NO: 1 or SEQ ID NO: 2 consisting of at least 15, preferably 20 to 30 or even more than 100 consecutive nucleotides is used as a probe to screen a DNA library for clones hybridizing to said fragment. The factors to be observed for hybridization are described in Sambrook et al, Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, chapters 9.47-9.57 and 11.45-11.49, 1989. Hybridizing clones are sequenced and DNA of clones comprising a complete coding region encoding a protein with more than 40% sequence identity to SEQ ID NO: 3 is purified. Said DNA can then be further processed by a number of routine recombinant DNA techniques such as restriction enzyme digestion, ligation, or polymerase chain reaction analysis.

The disclosure of SEQ ID NO: 1 enables a person skilled in the art to design oligonucleotides for polymerase chain reactions which attempt to amplify DNA fragments from templates comprising a sequence of nucleotides characterized by any continuous sequence of 15 and preferably 20 to 30 or more basepairs in SEQ ID NO: 1. Said nucleotides comprise a sequence of nucleotides which represents 15 and preferably 20 to 30 or more basepairs of SEQ ID NO: 1. Polymerase chain reactions performed using at least one such oligonucleotide and their amplification products constitute another embodiment of the present invention.

Knowing the nucleotide sequence of the *Arabidopsis* MKP1 gene and the amino acid sequence of the encoded protein it is possible to identify proteins interacting with AtMKP1 and to clone their corresponding genes using well known techniques. For example radioactively labeled AtMKP1 protein can be used for interactive cloning on cDNA expression libraries. AtMKP1 protein or parts thereof can be used to generate polyclonal or monoclonal antibodies specific for AtMKP1. The *AtMKP1* gene can be used to generate variants of AtMKP1 protein tagged with GST, MYK or His. Said antibodies and MKP1 variants allow to isolate native protein complexes by immunoprecipitation and to determine sequences of proteins present in these complexes by micro-sequencing. The resulting sequence information can in turn be used to clone corresponding genes. Alternatively, said antibodies or tagged MKP1 variants can be used to screen epitope libraries for epitopes which interact with AtMKP1 protein. The AtMKP1 protein and parts thereof, in particular the N-terminal 490 amino acid region and the C-terminal 492 amino acid region can also be used to search for interacting proteins with a Two-hybrid system (e.g. in yeast, in mammalian cells, or in bacteria). This allows to obtain sequence information about interacting proteins.

Based on the disclosed finding that AtMKP1 proteins are involved in a plant's abiotic environmental stress response, it becomes possible to engineer the corresponding signaling pathway, of which AtMKP1 is a part, to be chemically regulated due to chemical activation or repression of transgenes encoding AtMKP1 or proteins interacting therewith. Such plants can be obtained by transformation with the corresponding genes under control of chemically inducible promoters. Application of inducers is expected to modify the activity of the AtMKP1 signaling pathway and to result in altered adaptation to abiotic environmental stress. Alternatively, AtMKP1 protein or its interacting proteins can be used as targets for chemicals inhibiting or stimulating their activities which again is expected to modify abiotic stress responses.

EXAMPLES:

Example 1: *Cloning of the gene responsible for the mkp1 mutant phenotype*

Arabidopsis T-DNA insertion lines as produced by the INRA-Versailles and available from the Nottingham Arabidopsis Stock Center (NASC) are screened for sensitivity to methyl methanesulfonate (MMS) at a concentration of 100 ppm as described by Masson et al,

Genetics 146: 401-407, 1997. Plants which die in the presence of 100 ppm MMS are found in the family AAN4. Thus, the corresponding T-DNA insertion mutation is assumed to give rise to this hypersensitive phenotype. This assumption is supported by genetic analysis showing co-segregation of the hypersensitive phenotype with the T-DNA insertion.

Genomic DNA from the mutant plants is isolated as described by Dellaporta et al, Plant Mol Biol Reporter 1: 19-21, 1983. A fragment of genomic DNA flanking the right border of the inserted T-DNA is rescued essentially according to Bouché et al, Plant Mol Biol Reporter 14: 115-123, 1996, with minor modifications. Genomic DNA is digested with PstI, ethanol precipitated and resuspended in H₂O. DNA of vector pResc38 (Bouché et al supra) is digested with PstI and dephosphorylated with shrimp alkaline phosphatase. The phosphatase is heat inactivated, the vector DNA is ethanol precipitated and resuspended in H₂O. 2.5 µg of PstI digested genomic DNA and 2.5 µg of PstI digested and dephosphorylated vector are mixed and ligated overnight at room temperature in a total volume 100 µl in the presence of 10 units of T4 DNA ligase. The DNA of the ligation mixture is precipitated with ethanol, resuspended in 50 µl H₂O, and digested with XbaI in a total volume of 100 µl. XbaI digested DNA is precipitated with ethanol and resuspended in H₂O. A second overnight ligation reaction in the presence of 10 units T4 DNA ligase is performed in a total volume of 200 µl at room temperature to achieve circularization of DNA fragments. The DNA of the ligation mixture is again precipitated with ethanol, rinsed two times with 70% ethanol, dried and dissolved in 5 µl H₂O. Two 2 µl aliquots are used for electroporation of electrocompetent *E. coli* XL1-Blue cells (Stratagene) according to the manufacturer's instructions. Transformants containing the inserted T-DNA and adjacent *Arabidopsis* genomic DNA sequences are selected on plates with 50mg/l kanamycin. Single bacterial colonies are analyzed by isolation of plasmid DNA using QIAprep Spin Plasmid Kit (Qiagen) and restriction digestion with PstI and XbaI. Plasmid pBN1 containing 3.7 kb of inserted T-DNA linked to 5 kb of *Arabidopsis* DNA is identified. Sequencing of the junction site is performed using a primer directed towards the flanking plant DNA and having the nucleotide sequence 5' -GGTTCTACAGGACGTAACAAT-3' (SEQ ID NO: 14) complementary to T-DNA 41 nucleotides from the right border. Digestion of this clone with SstI allows isolation of a 960 bp fragment which when labelled with ³²P can be conveniently used as a probe to screen wild type *Arabidopsis* genomic and cDNA libraries in order to identify the wild type gene affected in the *mkp1* mutant line.

Example 2: Cloning of the *AtMKP1* wild-type gene

The 960 bp SstI fragment mentioned at the end of example 1 is labeled with ^{32}P by random oligonucleotide-primed synthesis (Feinberg et al, Anal Biochem 132: 6-13, 1983) for use as a probe in the following hybridization experiments.

Southern blot analysis of *Arabidopsis* wild type and *mkp1* DNA digested with EcoRV

confirms that in the *mkp1* genomic DNA the sequence hybridizing to the probe is linked to

T-DNA.

Northern blot analysis of *Arabidopsis* wild type RNA reveals the presence of a hybridizing transcript in RNA extracted from seven-day-old wild type seedlings. No such hybridizing fragment is detected in the corresponding RNA of *mkp1* seedlings.

A cDNA library (Elledge et al, 1991) and a genomic library (Stratagene) of wild type *Arabidopsis thaliana* ecotype Columbia is screened with the labelled SstI fragment

mentioned above. Screening of the bacteriophage λ libraries is performed according to the protocols described in chapter 6 of Ausubel et al, 1994, "Current protocols in molecular biology", John Wiley & Sons, Inc. Hybridization is performed as described by Church and Gilbert, Proc Natl Acad Sci USA 81: 1991-1995, 1984. Bacteriophage clones hybridizing to SstI fragment are subjected to *in vivo* excision of plasmids according to Elledge et al, Proc Natl Acad Sci USA 88: 1731-1735, 1991, and Stratagene protocols. Inserts of the plasmids obtained are further analyzed by sequencing.

By partial sequencing and alignment of ten overlapping clones (pBN5.1 to pBN5.10) isolated from the genomic library a continuous genomic sequence of 6356 bp (see SEQ ID NO: 1) is decoded.

Ten cDNA clones representing the same gene, one of them a 3.0 kb full-length cDNA (SEQ ID NO: 2), are isolated from the cDNA library.

Example 3: Sequence Analysis and Alignments

The 3 kb full-length cDNA clone of SEQ ID NO: 2 encodes an ORF with the start codon being defined by basepairs 298-300 and the stop codon by basepairs 2650-2652. The ORF encodes a protein consisting of 784 amino acids (SEQ ID NO: 3) and a predicted molecular mass of 86.0 kD. Alignment with the genomic sequence of SEQ ID NO: 1 reveals three introns. T-DNA is inserted within the coding sequence of the *mkp1* mutant DNA before basepair position 502 according to the numbering of SEQ ID NO: 2. The sub-sequence

VHCCQGVSR (SEQ ID NO: 4) found in SEQ ID NO: 3 can be interpreted as

corresponding to the mammalian sequence motif IHGXAGXXRS (SEQ ID NO: 5) defining

the family of protein tyrosine phosphatases, wherein the Ile at the first position can be

replaced by Val and the Ser at the last position can be replaced by Thr (Van Vactor et al,

Curr Opin Gen Dev 8: 112-126, 1998). Therefore it is concluded that the wild type ORF

encodes a protein tyrosine phosphatase with invariant aspartic acid, cysteine, and arginine

residues (Fauman et al, Trends Biochem Sci 21: 413-417, 1996) in positions 204, 235 and

241 according to SEQ ID NO: 3.

A data base search using the TFASTA program (Wisconsin Package Version 9.1, Genetics

Computer Group (GCG), Madison, Wisc.) reveals that the encoded phosphatase has a

significant similarity to dual specificity phosphatases. The closest homologue identified is

Xenopus laevis MAP kinase phosphatase (MKP; Lewis et al, J Cell Sci 108: 2885-2896,

1995) showing 38.1% identity and 52.5% similarity in a 140 amino acid overlap region. The

deduced AtMKP1 protein also has 36.0% identity and 52.5% similarity with a 140 amino

acid overlap region encoded by the rat 3CH134/CL100 cDNA representing a rat MKP1.

Essentially identical results are obtained when using the BLASTP 2.0.4 (Feb-24-1998)

program of the NCBI BLAST family of programs which, allowing gapped alignment,

compares an amino acid query sequence against a protein sequence database (Altschul et

al, Nucleic Acids Res. 25: 3389-3402, 1997). No higher plant homologues are identified.

The genomic position of the *AtMKP1* gene is determined by hybridization to filters containing genomic YAC clones publicly available from the Arabidopsis Biological Resource Center (Ohio, USA). The *AtMKP1* gene is found to map to chromosome 3 between markers ve022 and BGL1.

Example 4: *Complementation*

mkp1 mutant plants are transformed with DNA comprising the corresponding wild type

genomic DNA including promoter and polyadenylation signal to find out whether the cloned

wild type gene is able to complement the mutant *mkp1* phenotype.

mkp1 mutant plants harbor T-DNA containing the *NPTII* and *bar* marker genes under the

control of nos and CaMV35S promoters, respectively. Therefore, different marker genes are

used for the transformation construct. The vector used is a derivative of p1'barbi which is

highly efficient in *Arabidopsis* transformation (Mengiste et al, Plant J 12: 945-948, 1997). In

p1'barbi the EcoRI fragment containing 1'promoter, bar gene coding region, and CaMV 35S polyadenylation signal is inverted in respect to the T-DNA borders by EcoRI digestion and re-ligation. In the resulting plasmid the 1'promoter (Velten et al, EMBO J 3: 2723-2730, 1984) is directed towards the right border of the T-DNA. This plasmid is digested with BamHI and NheI, and the bar gene and CaMV 35S polyadenylation signal are replaced by a synthetic polylinker with the sites for the restriction enzymes BamHI, HpaI, ClaI, StuI and NheI. The resulting plasmid is digested with BamHI and HpaI and ligated to a BamHI-PvuII fragment of pROB1 (Bilang et al, Gene 100: 247-250, 1991) containing the hygromycin-B-resistance gene *hph* linked to the CaMV 35S polyadenylation signal. The T-DNA of the resulting binary vector p1'hygi contains the hygromycin resistance selectable marker gene under the control of the 1'promoter and unique cloning sites for the restriction enzymes *ClaI*, *StuI* and *NheI* located between the marker gene and the T-DNA right border. p1'hygi is used to insert the reconstructed *AtMKP1* gene as follows. Plasmid pBN1 of example 1 is digested with *PstI* and *MunI* and dephosphorylated. The restriction fragment containing the 3'portion of the *AtMKP1* gene and pBluescript-SK(+) is purified from the agarose gel and ligated to the *PstI*-*MunI* restriction fragment of the wild type genomic clone pBN5.2 (example 2) including the 5' end of the coding sequence of the *AtMKP1* gene and 2.4 kb of upstream sequences. The reconstructed *AtMKP1* gene is excised by *PstI* and *NotI* and after filling the ends is inserted into the *StuI* site of p1'hygi. The construct is introduced by transformation into *Agrobacterium tumefaciens* strain C58CIRif^R containing the non-oncogenic Ti plasmid pGV3101 (Van Larebeke et al, Nature 252: 169-170, 1974). T-DNA containing the reconstructed *AtMKP1* gene is transferred to mutant plants by the method of *in planta Agrobacterium* mediated gene transfer (Bechtold et al, C R Acad Sci Paris, Life Sci 316: 1194-1199, 1993). Seeds of infiltrated plants are grown on hygromycin-containing medium to screen for transformants. The progeny of selfed hygromycin resistant plants is analyzed for the segregation of hygromycin resistance. The families in which a 3:1 segregation ratio is observed are used to isolate homozygous lines bearing the newly introduced T-DNA inserted at a single genetic locus. The obtained hygromycin resistant lines are analyzed by Northern blot analysis for the restoration of *AtMKP1* expression. In these lines the restoration of transcription of the *AtMKP1* gene can be observed as well as the restoration of the wild type level of MMS resistance and ABA mediated stress responses. Complementation is not observed in plants transformed with p1'hygi only.

Example 5: Cloning of homologous sequences from other plant species

Use of *AtMKP1* cDNA as a probe for Southern hybridization with genomic DNA from other plant species such as *Sinapis alba* (mustard), *Lycopersicum esculentum* (tomato) and *Zea mays* (maize) is successful in the case of *Sinapis alba* which belongs to the same family as *Arabidopsis* (Brassicaceae).

Homologous sequences from the other species can be identified in a PCR approach using degenerate primers 1-3 below, wherein 1 is inosine, derived from the regions conserved between VH-PTP13 of *Clamydomonas eugametos* and *AtMKP1* protein:

Primer 1 (forward): 5' - AAY AAY GGI ATH ACI CAY ATH YT-3' (SEQ ID NO: 11);
 Primer 2 (reverse): 5' - YTG RCA IGC RAA ICC CAT RTT IGG-3' (SEQ ID NO: 12);
 Primer 3 (reverse): 5' - IGT CCA CAT IAR RTA IGC DAT IAC (SEQ ID NO: 13);

A PCR reaction is performed in a total volume of 50 µl containing 1x reaction buffer (Qiagen), 200 µM of each dNTP, 1.25 units of Taq polymerase (Qiagen), and 100 pmol of each primer. Reaction 1 is performed with primers 1 and 2 using genomic DNA from *Sinapis alba* (200 ng), *Lycopersicum esculentum* (400 ng), or *Zea mays* (600 ng) as the original template DNA. Amplification is carried out after an initial denaturation step of 3 min at 94°C, followed by 30 cycles of 30 sec at 94°C, 30 sec at 40°C, and 3 min at 72°C. The resulting amplification mixture is diluted 10³ fold. Reaction 2 is performed using 2 µl of the above dilution to provide the necessary template DNA. This time primers 1 and 3 are used under the same conditions as specified for reaction 1. The resulting amplification products are cloned into the T/A vector pCR2.1 (Invitrogen) and further analyzed by nucleotide sequencing.

Using this PCR approach it is possible to amplify sequences homologous to the *AtMKP1* gene from all the species mentioned above. Whereas the nucleotide sequence from *Sinapis alba* SaMKP1 (SEQ ID NO: 15 encoding SEQ ID NO: 16) is 90.8% identical to the *AtMKP1* sequence, the nucleotide sequence from *Lycopersicum esculentum* LeMKP1 (SEQ ID NO: 17 encoding SEQ ID NO: 18) is 72.3% and the *Zea mays* sequence

ZmMKP1 (SEQ ID NO: 19 encoding SEQ ID NO: 20) 71.8% identical. The fragments hybridize to genomic DNA from corresponding species under the usual hybridization conditions for Southern blot analysis. The fragments can be used as probes to screen cDNA libraries for corresponding cDNA sequences.

The 243 bp *ZmMKP1* fragment amplifying from maize DNA is used as a probe to screen a maize cDNA library (Clontech) made in the Lambda ZAP® II Vector (Clontech) from "Blizzard" hybrid etiolated shoots, which were treated with the herbicide safener Benoxacor. The titer of the library is determined as 3×10^9 pfu/ml.

Library screening is conducted as described in the Clontech Lambda Library Protocol Handbook, with some slight modifications. Briefly, a single colony of XL-1 Blue is picked and incubated overnight at 37°C in LB medium, containing 10mM $MgSO_4$ and 0.2% maltose. 600 µl of stationary phase grown bacteria for each 150 mm plate is combined with 100 µl of phage library dilution in sterile 1x lambda dilution buffer (100mM NaCl; 10mM $MgSO_4$; 35mM Tris-HCl, pH7.5) to yield approximately 30,000 pfu per plate. This mixture is incubated at 37°C for 15 minutes, subsequently 7 ml of melted LB soft top agarose (at 48°C) is added to the cell suspension for each 150 mm plate, shortly mixed and then poured on two-day-old LB $_{MgSO_4}$ agar plates, which have been pre-warmed to 37°C for four hours. The plates are then incubated at 37°C until plaques reach appropriate sizes (after about 8 to 9 hours). After chilling the plates at 4°C for one hour, phage particles are transferred to Hybond N nitrocellulose filters and the orientation of each filter to its plate is recorded with a waterproof pen. The filters are then treated by placing them on Whatman 3MM paper saturated with the appropriate solution. The treatments include denaturation solution (0.5M NaOH; 1.5M NaCl) for 2 minutes, followed by neutralization solution (0.5M Tris-HCl, pH 7.2; 1.5M NaCl; 1mM EDTA) for 3 minutes and 2x SSC for 3 minutes. DNA is subsequently crosslinked to the filters by UV.

Filters are then pre-hybridized, hybridized with the radioactively labeled *ZmMKP1* fragment and washed as described in Sambrook et al, Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, chapters 9.47-9.57 and 11.45-11.49, 1989. An agar plug from the position of a positive plaque is then removed from the master plate and incubated overnight at 4°C in 1 ml of 1x lambda dilution buffer, containing 20 µl of chloroform. Each titer is determined and the phages are re-plated to obtain approximately 200 to 500 plaques on a 150 mm plate for a secondary screen as described above. Single plaques of interest

are collected from the agar plates and incubated over night at 4°C in 500 µl of 1× lambda dilution buffer and 20 µl of chloroform.

The pBluescript phagemid is excised from the λZAP™ vector as described by the *In Vivo* Excision Protocol using the EXAssis/SOLR System in the Stratagene Uni-ZAP™ XR Library Instruction Manual (1993). A 1/100 dilution is made of XL1-Blue MRF' and SOLR overnight cultures (at 30°C) and incubated at 37°C for 2-3 hours. XL1-Blue MRF' cells are then pelleted for 10 minutes at 1,500×g and re-suspended at an $OD_{600} = 1.0$ in 10mM $MgSO_4$. 200 µl of these XL1-Blue cells are then combined with 250 µl of phage stock and 1 µl of EXAssist helper phage in a 50 ml conical tube and incubated at 37°C for 15 minutes. 3 ml of LB broth is added and incubated at 37°C for 5 hours, after which the cells are pelleted for 15 minutes at 2,000×g and the supernatant transferred to a new tube. The tube is then heated at 70°C for 15 minutes and centrifuged again for 15 minutes at 4,000×g. The supernatant, containing the excised phagemid pBluescript packaged as filamentous phage particles, is decanted into a new tube. 10 and 100 µl of this phage stock are then added to two tubes with 200 µl of SOLR cells that have been allowed to grow to $OD_{600} = 0.5-1.0$ before being removed from the incubator and further incubated at room temperature. The tubes are incubated at 37°C for 15 minutes, followed by plating 10-50 µl from each tube on LB^{amp} (50 µg/ml) and over night incubation at 37°C.

The positive clones are checked for insert size by *EcoRI/XhoI* double digestion and end-sequencing with T3 and T7 promoter primers (Promega).

Screening of 360,000 pfu of the library results in three identical clones of 2.2 kb containing the 3' poly(A) tail, but lacking part of the 5' end, including the translation initiation site. The gene corresponding to the identified partial cDNA clone is named *ZmMKP2*, as it is not identical with the *ZmMKP1* fragment used as the probe (92.3% identity on the nucleotide level over the 196 bp fragment flanked by the primers 1 and 3). An additional 213 nucleotides are amplified and cloned by 5' RACE (rapid amplification of cDNA ends) carried out following the instructions of the 5'/3' RACE Kit (Boehringer Mannheim) resulting in a longer cDNA sequence of 2,452 bp but still not complete, judged by the predicted mRNA length from the RNA gel blot analysis and the absence of a possible translation initiation site.

The sequence information gained from the *ZmMKP2* cDNA including the additional 213 nucleotides obtained by 5' RACE (SEQ ID NO: 21 encoding SEQ ID NO: 22) is used to

design two additional backward oriented degenerate primers wherein 1 is inosine to 3' regions conserved between the deduced peptide sequences of *ZmMKP2* and *AtMKP1*:

Primer 4 (reverse): 5'-GCI GCY TTI GCR TCY TTY TCC-3' (SEQ ID NO: 25);

Primer 5 (reverse): 5'-YTC ICK IGC IGG IAR RTG IGT YTC-3' (SEQ ID NO: 26)

These primers are used to PCR amplify a larger fragment of a MAP kinase phosphatase gene from tomato. The amplified and cloned 522 bp long fragment is not identical to

LeMKP1. Therefore, its corresponding gene is named *LeMKP2* (SEQ ID NO: 23 encoding SEQ ID NO: 24; 75% identity on the nucleotide level over the stretch of 196 bp of

ZmMKP1 flanked by primers 1 and 3). The origins of all identified MAP kinase phosphatase homologous gene sequences are confirmed by Southern blot analysis.

The following Table shows an alignment of a continuous stretch of 312 amino acids of *AtMKP1* with the related amino acid sequence of *ZmMKP2*.

139	AtMKP1	KREKIAFTDKESKVAHDHIVGDAVAKDKSIKNGITTHIINCVGFI	188
24	ZmMKP2	RKDQIAFTDKESKVAHDHVLGDAVAKNRDILRNKGITTHVINCVG	73
189	AtMKP1	EYFKSDFCYRSIWLQDSPSEDITSLYDFVDFEVRQSGRI	238
74	ZmMKP2	EYFKSDFCYRSIWLQDSPSEDITSLYDFVDFEVRQSGRI	123
239	AtMKP1	VSRSTSLVLAFLMWRBQSFDDAFQYVKSARGIADFNMGFA	288
124	ZmMKP2	VSRSTSLVLAFLMWRBQSFDDAFQYVKSARGIADFNMGFA	173
289	AtMKP1	VHAFPLSPSTSLRMKMSFHSFYDPLHLVFKLNDPCPSL	338
174	ZmMKP2	VHAFPLSPNSVLRMYRMAFHSQYAPLHLVFKLNDPSPATL	223
339	AtMKP1	LPSAIYTWVGRQCEITMEKDAKAAVCQIARYEKVEAFIM	388
224	ZmMKP2	VLSSLYVWVGAKCDPVMKKDAKAAAFQVVRKYKVGHTK	273
389	AtMKP1	WDAFASITLPMIGG.....SVTKVQPGDKKVDAYNLD	428
274	ZmMKP2	WDAFSSMPFNSDSNTKISKDQIDSAKSDPCSRKNESYDA	323
429	AtMKP1	GGFVPTLASNNHEHTLPARE	450
324	ZmMKP2	GGVVPASTSGAGDEHTLPARE	345

SEQUENCE LISTING

<110> Novartis AG

<120> Organic compounds

<130> Silencing Gene

<140>

<141>

<160> 26

<170> PatentIn Ver. 2.1

<210> 1

<211> 6356

<212> DNA

<213> Arabidopsis thaliana

<400> 1

gttcaagggtg gttgtatttcn cagtattaga aagaggctt ctagaggag ttcttaatt 60
atatctgaa caggccgga ttggtgtct tagattaaag tgaatgctt agatctggtg 120
acgatggtg tagagatat atagtamaat gatagaacat cagaataact ataagtcacc 180
atatcttaa aaaataaatg ctatagaatt ttgttttg taattgttat aactctacaa 240
agttatgtgt tatagagttt tactagtttc atcgttatca tgtgtatagt ataaacaa 300
aaaagaatt ttaatatct gaaaacataa aatttaata aatgtatgtg agtataagaa 360
aaaagaagaa aagaacaa cgtaaaaaat aaaaatcat tatacatat acaatttca 420
aaagatcaat gttaaattt attagtctt ttcttatgt ttcatgcaaa tgcataagta 480
tcaacttct ttaactaaag attagtgtt gcttttaagc aagaaactat caaagagtca 540
aagacattga tgtaaactt agagaacggg atcttctac aatcttat ttctccctt 600
ttttttaca acgaacgata gtttagtaagc atgaaatcgg ttcaaatgtt ggtatcaggt 660
gggcataatc ctggttcggt aaaaagaaac aataataat atgttgagc tagagagaag 720
aagctcatag ttctgttag aaaaacgtg gagatlgag agaatatca aagagagcga 780
agagaagagt aaacctagaa gatacaaaag acgctatcaa aagatgtt tatgtgtta 840
tgaacatct acttagcca gtaataat taaatgtcat tttygaaag aaaaaaaat 900
ggtgtcattt ttaggatat aatatctt caacaagatg tttagact tcaacataa 960
gatactttaa aactacataa ttatctct cgttaaaac ttacatcaa caaaagta 1020
aatctgacaa gaagagttt tcaaatagtt atactgtatg tgtttattt attcacatga 1080
aaagtgtcaa acaacactg ggtacatat ttcatgctt aaatcgtat aacaacatt 1140
aagttaatca gtaatttga tgcctaaact aaaaatgtg accatttga ccaatgagca 1200
ttttttgaa ataaactac tgcatacgac ctggtcaaac acatgaaact ccaactgctt 1260
taaaactacc tttttaacc ttgacgtcc attttatc actaatatc atgttaatca 1320
aaccaactaa attatgattc gaaatttcga gtagatgtg ttagaagtg tgtgaatgat 1380
aaccaactaa attatgattc gaaatttcga gtagatgtg ttagaagtg tgtgaatgat 1440
tcgttccaaa acatgtatga taaaatgttaa ttacatatcaa aaacttaag ccagtatga 1500
aaataaaaaa aataaataag tataatgttc atatatgtc ccaaccataa tttrgtttg 1560
atgaggaaaa gttctttt tttrttct tttrttct tagttaaag acataaaaaa 1620
aatataatt acaaacaaag caaaaaaaa gaataaataa aacaacaaatc attaactgca 1680
aatcaattat cagagggaaa agttataaag aaatgtcaac caatgagtat catatcatc 1740
atgcttttt agaccctct tttaattca tcaagattta gtttgttta taaatttga 1800
gtaaaattct aaatgaatca atctacaat tttrttccca catratgta acgataaaat 1860
ttaagatcaa caattactc gttaatrttt tttrttct tttrttctgataa aaatatttga 1920
attatgatat attagcatg tttrttgtratt gttatgtatg taaatattc tagttcaaaa 1980

- What is claimed is:**
1. A DNA comprising an open reading frame encoding a protein characterized by a component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO: 3
 2. The DNA according to claim 1 comprising an open reading frame encoding a plant MAP kinase phosphatase.
 3. The DNA according to claim 1 wherein the open reading frame encodes an amino acid sequence selected from the group of amino acid sequences described in SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.
 4. The DNA according to claim 1, wherein the open reading frame encodes a protein characterized by the amino acid sequence of SEQ ID NO: 3
 5. The DNA according to claim 1 characterized by the nucleotide sequence of SEQ ID NO: 1
 6. The DNA according to claim 1 wherein the open reading frame encodes a protein contributing to repair of DNA damage in a plant cell.
 7. The DNA according to claim 1 wherein the open reading frame encodes a protein conferring hypersensitivity to treatment with methyl methanesulfonate (MMS).
 8. The DNA according to claim 7 wherein the open reading frame encodes a protein conferring hypersensitivity to treatment with UV light or X-rays.
 9. The DNA according to claim 7 wherein the open reading frame encodes a protein interfering with abscisic acid signal transduction.
 10. The protein encoded by the open reading frame of any one of claims 1 to 9.
 11. A method of producing DNA according to claim 1, comprising
 - screening a DNA library for clones which are capable of hybridizing to a fragment of the DNA defined by SEQ ID NO: 1, wherein said fragment has a length of at least 15 nucleotides;
 - sequencing hybridizing clones;
 - purifying vector DNA of clones comprising an open reading frame encoding a protein with more than 40% sequence identity to SEQ ID NO: 3
 - optionally further processing the purified DNA.
 12. A polymerase chain reaction wherein at least one oligonucleotide used comprises a sequence of nucleotides which represents 15 or more basepairs of SEQ ID NO: 1.

gaataataaa tggtrtctat agcatgaatt taataaataa ataagactga ataacatag 2040
 gtaataaact ttgttctt ttgtaaatgt aaaaatagaa aaaaatagaa 2100
 ataatataa ttacattcaa ttgtaagaatt aaaaatagaa gtaataaact 2160
 agaaaataaa gaattataa tcttgtaaga gcaataaag aaaaatagaa 2220
 ttcaagaatt tcaacgctt tcttaataa agaatccaat ttgtaagaatt 2280
 ataataataa taataataa aatcgttaac acttaataa actccaataa 2340
 cgaattataa ttatgttaat ttgttaataa ttgtttcaat ttctcagtt 2400
 atctctctt tcttttgt ttctctctt tctcgtctt gtttcccaa tctctctt 2460
 ctctctctt gaagaaataa aaataaataa tctaaacttg acgctctctt taatcttaac 2520
 caatcgttaa gttccaataa tctctctt actcctaat actcctaat 2580
 ccaaggaatt tgcctcaacc ccttttgtg tttrtaataa tttrtaataa 2640
 ctgttctcgt ctctagat ttggtttca ttctgttt ttctgttt 2700
 ttcatrttgt gtttctgagt ttatctagtt agctgaatt tttaattctg 2760
 gctttctct cttrctag ttgcttaatt cgtcttgat ctgattcgtg ttatctct 2820
 ttccctctgt tttagtttt tttaggttt tttaggttt tttaggttt 2880
 gtttatrttt agatcagat ttgatataa gcttctcgtg aactaaag ttctaggaag 2940
 aagcaagaac cttgtttta caattgttt ttgttaggaat ttgtctgatt ttggtttgata 3000
 aagttgaatt cttrgagt tttaatttt gaataatttt gatgaatttt agagaggttt 3060
 cgttggttaa ttgattgaatt cctccttgt tttaagaaat gtttttgtg ttctcctt 3120
 gtttctgttt aggtactga ttcaagtttt ttgaagtttt ttgaagtttt 3180
 cgttgcttat ttgttctggt ttgttctggt ttgttctggt ttgttctggt 3240
 caatctataa cagcaaggt ttgttttgt ttgttttgt ttgttttgt 3300
 gtaggagctt agatcagtt tttaaggttt tttaaggttt tttaaggttt 3360
 ccaataacc ttgcttggttt ttgcttggttt ttgcttggttt ttgcttggttt 3420
 agtagcgttt ttgcttggttt ttgcttggttt ttgcttggttt ttgcttggttt 3480
 ataaagatt ttcaaggttt ttgcttggttt ttgcttggttt ttgcttggttt 3540
 acagagctt ttgcttggttt ttgcttggttt ttgcttggttt ttgcttggttt 3600
 gttcgttaa ttcaagtttt ttgcttggttt ttgcttggttt ttgcttggttt 3660
 cagtaggttt agcttagtt ttgcttggttt ttgcttggttt ttgcttggttt 3720
 aaagtgttaa gattcttttt ttgcttggttt ttgcttggttt ttgcttggttt 3780
 ttgcttatct gatgttggttt ttgcttggttt ttgcttggttt ttgcttggttt 3840
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 3900
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 3960
 caatctcttt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4020
 gttctctggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4080
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4140
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4200
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4260
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4320
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4380
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4440
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4500
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4560
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4620
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4680
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4740
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4800
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4860
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4920
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 4980
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 5040
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 5100
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 5160
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 5220
 ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt ttgcttaggt 5280

tacaatrg tgaatccat tigtctaga tgaactgac taattac taattgtaa gtcgtgac 5340
 taataatagc taaacttg gaggctctga tactttac taattgtaa gtcgtgac 5400
 tcatcttctg catgtcttat ccaaccatc tatattcga agatgaaat tacaatcga 5460
 gcaattcat tactgactac tgaagacgt taggtagaat cctatgatt tcaagcattg 5520
 tatgtatttg ttatctct agtggtctg atgttccaa ctgtttatga tcttttgt 5580
 gtttgttaact gataagtgc ttctctctc tgttaacaga tagttagga atctgagat 5640
 cagaagagt tgcagacat gctgagcgc ctataacac caccgcaag ctctacat 5700
 ttaacttgt ttctttcac agatctctc aaccgcaaca ctttccat ttcaagcaga 5760
 gtaattcat agctcaggt agaatctc gaaagcagtc tgtaacat catctaca 5820
 gttgcatcgg aatacaatcg ttagtctg atattgta atgtctat gatcatgaat 5880
 ttgaattaga ggaattgga acaaaaaa cttagagct cgaatgacg gttttaca 5940
 aattctcata gacctatt gattcttg attactct gttgcaagg tctctgtc 6000
 tatggaagt gatgtggg gaaacact tctgtacag tgggaaaaa actctctc 6060
 ctctctca tcaatgaaa atctcaggg gcaattata gtaatcag atataaat 6120
 tgaagtta gggtctat gaggatttg atgactgt tacaattgt acatataca 6180
 tcagcagcac aatagatt ttgtaaact acatgttat caagtaaaag tactatgag 6240
 atgttgaagt ctatgaa attagtaa atagttctc ttctttat tctctcaa aacag 6356

<210> 2

<211> 3059

<212> DNA

<213> Arabidopsis thaliana

<400> 2

ctctctcc gcttcttc ccaatctc ctctctc tctctgaaga aaataata 60
 aagaattca cttgacgc tcttaatc ttaactac cgttttaag tctgaataga 120
 ctgattgag ctgattgt atgttagat caatgtga tttaagct tgcctgaat 180
 acaagtca tgaagagc aagaacct gtttacct gtaattgt agaatctgt 240
 ctgatttg gtttaagg tgaagctt gatttgtaa tttagata agattgag 300
 gtgggaag agtatgcgt gggaattg gaagctct ctgttcaa gaaatctt 360
 tggcgtct cctctgtc tgcctcag actgcatc aagtctga ggttgatga 420
 caagcctga acatcctg tgcattagt tctggcga gtcgaagat tccagctc 480
 ccttgacac ctgttcaa tcaaacag aagctagag ctgttgc accattgcag 540
 cctctgca ttctagag gacttagc gattgctta agcggttc ggaatgattc 600
 gttgaattgc ctatccac aacactag ggaacaaa cggggagag atgaaactc 660
 gatctcat caagcagc gctgttaaca gataagct ctgtctag taagaggaag 720
 aagattct tcttgata agaatgtca aagttctg atcatata tgtgggtga 780
 gatgctgt cgaagaca gacatact aaaaacatg gaatcagca tatctgaac 840
 tgcgttgt ttatctgc ggaattct aagctgat ttgttaag atcttgtg 900
 ttaagata gtcgtcaga ggaatcaat agtatctgt acgatgtgt tgaacttt 960
 gaagattga gggagcaag tggaggaat ttgttcat gttgtcaag gtttcaaga 1020
 tcaactct tggtaatag atacttgat tgaagagag gccaagtct tgaatgaga 1080
 ttcaatag tgaagtct tagagttat gctgactca acatgggt tgcctgcaa 1140
 ttgttaaat gcaaaagag gttcactag tctccgcta gccctacc ctactaga 1200
 atgtacaaa tgcctcaca ctctctat gaccttgc atcttgtc aaaaactgtg 1260
 aatgaatcat gcccggtc tctgattca agagtgcat ttatctca gttacctc 1320
 gcaattaca ttgggttg taggcagt gaaacctca tggagaaga tgcataagct 1380
 gctgttgt agatgtcg atatgagaa gtcgaagca ctattatgt gttcagaga 1440
 ggtgaattag ctgtttat ctgggatga ttgcaagca ttgtccat gattggggc 1500
 tctgttaat aagttcaac agttgacag aagtcgag catataatc ggaatttgag 1560
 attttcaga aagccatga gggaggttt gtcgaactc tagcatc caacaacga 1620
 catgagact atctctgc aagggaatc agttgagct cactgaatg taagtctga 1680
 tcaaggttg acaagggtt tctgtatgc tccaaaacgc cactatcag gttctatca 1740
 gatcgtga tgaatcgtc atatgagaa gtcgaagca ttgtccat gattggggc 1800

actgcgtcgc ctccttct ctcctccgat tctgtatgt caacaattc aggaatagc 1860
ttaaagatt tctctcaatc ctcgtgaagt tctgtcctga gacctctat tccaaccag 1920
ctgacattgc ctaatttc cagctatcc ctcctccct cccaactc tctaagaa 1980
tctcgttgtg tcaattatc tctcaaccg tccaataa gaaagctc acctctct 2040
gctgaagcta gaggcagct gaaagatc ctgaagtgc caggttgc tgaatccaac 2100
agaggcacac ctgctttac ttacatccg gatgatagt atgacatagt ctcaatcg 2160
gaggtatta gaaaggcga tctatatca ccaagtgtt gcaaggagc aagtgtagat 2220
tcgatttgc cagaagaaga aatatatcc ttaatcagt gcagttaac tgacagacac 2280
aaatcgggag gtgatactga tagctctgc cagccttag catgtcgtg gccaaagt 2340
gagatgatta caaactgag cagagctac ttagatcag aatcgttat agcaatccg 2400
tgccaagt atgctgtag gaaacgggt agtaggaat tgaatattg gatcggaaag 2460
tcatctctc ttgatacaaa ctgtctcta gtatagca acaagcggc agacactgtg 2520
gagaatgtg attggtaca aatggtga tccatttgt gtcaagtga ctgccaataa 2580
gatacccta taaagatagt taggaaatc gagatcaga cagattgtc agcactgtg 2640
agcgctat aacaccacc cgcaagctt acacattac tctgtttt ttccacagat 2700
tcttcaacc gcaacactt tccatttca gacagatat tcatcagct caggaactc 2760
tgtgttag gagtgtag ttggggaaa caactctct gtacagtggt gaaaaact 2820
ctctcttc ttctatcac atgaaatcc tcaaggcca ttaatagat gatcagatta 2880
taaatgta agtttagg cttrtagg attttagg acttgta atgtttacat 2940
atacactag cagacaata gatrttgtt aaactacat gtatcaag taagaact 3000
atgtagatg tgaaagtcaa ttgaagaat agttaatgat aaaaaaaaaa 3059

<210> 3

<211> 784

<212> PRT

<213> Arabidopsis thaliana

<400> 3

Met Val Gly Arg Glu Asp Ala Met Gly Asn Asp Glu Ala Pro Gly 1
5
10
15

Ser Lys Lys Met Phe Trp Arg Ser Ala Ser Trp Ser Ala Ser Arg Thr 20
25
30

Ala Ser Glu Val Pro Glu Gly Asp Glu Glu Ser Leu Asn Ile Pro Cys 35
40
45

Ala Ile Ser Ser Gly Pro Ser Arg Arg Cys Pro Ala Ala Pro Leu Thr 50
55
60

Pro Arg Ser His His Asn Ser Lys Ala Arg Ala Cys Leu Pro Pro Leu 65
70
75
80

Gln Pro Leu Ala Ile Ser Arg Arg Ser Leu Asp Glu Trp Pro Lys Ala 85
90
95

Gly Ser Asp Asp Val Gly Glu Trp Pro His Pro Pro Thr Pro Ser Gly 100
105
110

Asn Lys Thr Gly Glu Arg Leu Lys Leu Asp Ser Ser Thr Gln Gln 115
120
125

Arg Val Thr Asp Lys Ser Ser Gly Leu Ala Lys Arg Glu Lys Ile Ala 130
135
140

Phe Phe Asp Lys Gln Cys Ser Lys Val Ala Asp His Ile Tyr Val Gly 145
 150 155
 Gly Asp Ala Val Ala Lys Asp Lys Ser Ile Leu Lys Asn Asn Gly Ile 165
 170 175
 Thr His Ile Leu Asn Cys Val Gly Phe Ile Cys Pro Gln Tyr Phe Lys 180
 185
 Ser Asp Phe Cys Tyr Arg Ser Leu Trp Leu Gln Asp Ser Pro Ser Gln 195
 200 205
 Asp Ile Thr Ser Ile Leu Tyr Asp Val Phe Asp Tyr Phe Gln Asp Val 210
 215
 Arg Gln Gln Ser Gly Arg Ile Phe Val His Cys Cys Gln Gly Val Ser 225
 230 235
 Arg Ser Thr Ser Leu Val Ile Ala Tyr Leu Met Trp Arg Gln Gly Gln 245
 250 255
 Ser Phe Asp Asp Ala Phe Gln Tyr Val Lys Ser Ala Arg Gly Ile Ala 260
 265
 Asp Pro Asn Met Gly Phe Ala Cys Gln Leu Leu Gln Cys Gln Lys Arg 275
 280 285
 Val His Ala Phe Pro Leu Ser Pro Thr Ser Leu Leu Arg Met Tyr Lys 290
 295 300
 Met Ser Pro His Ser Pro Tyr Asp Pro Leu His Leu Val Pro Lys Leu 305
 310 315
 Leu Asn Asp Pro Cys Pro Gly Ser Leu Asp Ser Arg Gly Ala Phe Ile 325
 330 335
 Ile Gln Leu Pro Ser Ala Ile Tyr Ile Trp Val Gly Arg Gln Cys Gln 340
 345
 Thr Ile Met Gln Lys Asp Ala Lys Ala Val Cys Gln Ile Ala Arg 355
 360 365
 Tyr Gln Lys Val Gln Ala Pro Ile Met Val Val Arg Gln Gly Asp Gln 370
 375 380
 Pro Val Tyr Tyr Trp Asp Ala Phe Ala Ser Ile Leu Pro Met Ile Gly 385
 390 395
 Gly Ser Val Ile Lys Val Gln Pro Gly Asp Arg Lys Val Asp Ala Tyr 405
 410 415
 Asn Leu Asp Phe Gln Ile Phe Gln Lys Ala Ile Gln Gly Gly Phe Val 420
 425 430
 Pro Thr Leu Ala Ser Ser Asn Asn Gln His Gln Thr His Leu Pro Ala

Arg Glu Asn Ser Trp Ser Ser Leu Lys Cys Lys Phe Ala Ser Arg Phe 445
Asp Lys Gly Phe Arg Tyr Val Ser Lys Thr Pro Leu Ser Arg Val Tyr 465
Ser Asp Ser Met Ile Val His Ser Ser Gly Ser Pro Phe Leu Ser Pro Asp Ser 470
Thr Ser Ser Ser Thr Ala Ser Pro Pro Phe Leu Ser Phe Ser Gln Ser 485
Ser Asp Ser Met Ile Val His Ser Ser Gly Ser Pro Phe Leu Ser Arg Val Tyr 490
Thr Ser Ser Ser Thr Ala Ser Pro Pro Phe Leu Ser Phe Ser Gln Ser 500
Val Cys Ser Thr Asn Ser Gly Asn Ser Leu Lys Ser Phe Ser Gln Ser 510
Ser Gly Arg Ser Ser Leu Arg Pro Ser Ile Pro Ser Leu Thr Leu 520
Ser Gly Arg Ser Ser Leu Arg Pro Ser Ile Pro Ser Leu Thr Leu 535
Ile Leu Ser Ser Leu Ser Leu Ser Leu Ser Leu Ser Leu Thr Leu 540
Cys Ser Thr Val Asn Thr Phe Leu Gln Pro Ser Gln Thr Ser Pro Lys 550
Glu Ser Arg Gly Val Asn Thr Phe Leu Gln Pro Ser Gln Thr Ser Pro Lys 555
Ala Ser Pro Ser Leu Ala Glu Arg Arg Gly Ser Leu Lys Gly Ser Leu 565
Lys Leu Pro Gly Leu Ala Asp Ser Asn Arg Gly Thr Pro Ala Phe Thr 570
Lys Leu Pro Gly Leu Ala Asp Ser Asn Arg Gly Thr Pro Ala Phe Thr 575
Lys Leu Pro Gly Leu Ala Asp Ser Asn Arg Gly Thr Pro Ala Phe Thr 580
Lys Leu Pro Gly Leu Ala Asp Ser Asn Arg Gly Thr Pro Ala Phe Thr 585
Lys Leu Pro Gly Leu Ala Asp Ser Asn Arg Gly Thr Pro Ala Phe Thr 590
Lys Leu Pro Gly Leu Ala Asp Ser Asn Arg Gly Thr Pro Ala Phe Thr 600
Leu His Pro Asp Asp Ser Asn Asp Ile Val Phe Asn Leu Glu Gly Ile 605
Arg Asn Gly Asp Leu Tyr Pro Pro Ser Asp Cys Lys Gly Thr Ser Val 615
Asp Ser Asp Leu Pro Glu Lys Glu Ile Ile Ser Leu Ile Ser Cys Ser 620
Asp Ser Asp Leu Pro Glu Lys Glu Ile Ile Ser Leu Ile Ser Cys Ser 630
Lys Ser Asp Arg His Lys Ser Gly Asp Thr Asp Ser Ser Gly Gln 635
Pro Leu Ala Cys Arg Trp Pro Ser Met Glu Met Ile Thr Lys Leu Ser 640
Arg Ala Tyr Leu Asp Ser Glu Ser Val Ile Ala Ile Pro Leu Pro Ser 645
Asp Ala Val Gly Glu Thr Gly Ser Arg Asn Leu Tyr Ile Trp Ile Gly 650
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 655
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 660
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 665
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 670
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 675
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 680
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 685
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 690
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 695
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 700
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 705
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 710
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 715
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 720
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 725
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 730
Lys Ser Phe Ser Leu Asp Asn Cys Ser Leu Val Asp Ser Asn Lys 735

<210> 7
<211> 12
<212> PRT
<213> Arabidopsis thaliana
<400> 7
Phe Val His Cys Cys Gln Gly Val Ser Arg Ser Thr

<210> 6
<211> 14
<212> PRT
<213> Arabidopsis thaliana
<400> 6
Thr Ser Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val
1
5
10

<210> 5
<211> 10
<212> PRT
<213> Artificial Sequence
<220> Description of Artificial Sequence: mammalian
<223> sequence motif defining the family of PTPs
<400> 5
Ile His Cys Xaa Ala Gly Xaa Xaa Arg Ser
1
5
10

<210> 4
<211> 10
<212> PRT
<213> Arabidopsis thaliana
<400> 4
Val His Cys Cys Gln Gly Val Ser Arg Ser
1
5
10

Ala Ala Asp Thr Val Glu Asn Val Asp Trp Val Gln Ile Gly Glu Ser 740
Ile Leu Cys Gln Met Asp Leu Pro Lys Asp Thr Ile Lys Ile Val 745
Arg Glu Ser Glu Asp Gln Thr Glu Leu Leu Ala Leu Ser Ala Leu 750
755
760
765
770
775
780

1 5 10

<210> 8
<211> 4
<212> PRT
<213> Arabidopsis thaliana
<400> 8
Phe Val His Cys
1

<210> 9
<211> 6
<212> PRT
<213> Arabidopsis thaliana
<400> 9
Gln Gly Val Ser Arg Ser
1
5

<210> 10
<211> 5
<212> PRT
<213> Arabidopsis thaliana
<400> 10
Tyr Phe Lys Ser Asp
1
5

<210> 11
<211> 23
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: degenerate
primer

<400> 11
aayaaygna thacncayat hyt

<210> 12
<211> 24
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: degenerate
primer
<400> 12

24

ytgcanger aannccatt tngg

<210> 13

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: degenerate primer

<400> 13

ngtccacatn arrtngcda tmac

24

<210> 14

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 14

gatttccaca gacgtaaca t

21

<210> 15

<211> 196

<212> DNA

<213> Synapis alba

<400> 15

caactgcgtt ggttcact gtccgaata ttccaagctt gatttttgtt accgtcgtt 60
gtgttcagt gatagtccat cagaggatat aactagcat ctctacgatg tctttgacta 120
cttgaagac gttaggagc aaagtggag gatttttgtt cactgtgtc aagcggttc 180
acgtccacc tccctg 196

<210> 16

<211> 65

<212> PRT

<213> Synapis alba

<400> 16

Asn Cys Val Gly Phe Ile Cys Pro Glu Tyr Phe Lys Ser Asp Phe Cys
1 5 10 15

Tyr Arg Ser Leu Trp Leu Arg Asp Ser Pro Ser Glu Asp Ile Thr Ser
20 25 30

Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val Arg Glu Glu Ser
35 40 45

Gly Arg Ile Phe Val His Cys Cys Glu Gly Val Ser Arg Ser Thr Ser
50 55 60

Leu

65

<210> 17
<211> 196
<212> DNA
<213> Lycopersicon esculentum

<400> 17
aaactgtta ggttagt gtcctaata ctttaagat gacctgtat acaagacact 60
ttggctgag gatagccca ctgaggacat caaagatat cttatgat tcttgatta 120
cttgaagat gtcatgaac aagtgaggag tgccttgta cactgctcc aggggtgtc 180
ccgatagcc tccctg 196

<210> 18

<211> 65
<212> PRT
<213> Lycopersicon esculentum

<400> 18
Asn Cys Val Gly Phe Ser Cys Pro Gln Tyr Phe Lys Asp Leu Val 1
5
Tyr Lys Thr Leu Trp Leu Gln Asp Ser Pro Thr Gln Asp Ile Thr Ser 20
30
Ile Leu Tyr Asp Val Phe Asp Tyr Phe Gln Asp Val His Gln Gln Gly 35
40
Gly Ser Val Phe Val His Cys Phe Gln Gly Val Ser Arg Ser Ala Ser 50
60

Leu
65

<210> 19
<211> 196
<212> DNA
<213> Zea mays

<400> 19
caactgcatg ggttcgtct gcccgaaata cttcaagctt gacctgtct accgacacct 60
ctgctgcag gacagcccca ccgaggacac caccagcatc cttacgacg tgttgatta 120
ctttaggac gtcaaggagc aggtggccg cgtgttgctg catgctgcc aggggtgtc 180
ggctccacg cctctg 196

<210> 20

<211> 65
<212> PRT
<213> Zea mays

<400> 20
Asn Cys Met Gly Phe Val Cys Pro Lys Tyr Phe Lys Ser Asp Leu Val 1
5
10
15

Tyr Arg Thr Leu Trp Leu Gln Asp Ser Pro Thr Gln Asp Thr Thr Ser	20	25	30
Ile Leu Tyr Asp Val Phe Asp Tyr Phe Gln Asp Val Arg Gln Gly	35	40	45
Gly Arg Val Phe Val His Cys Cys Gln Gly Val Ser Arg Ser Thr Pro	50	55	60
	Leu	65	

```
<210> 21
<211> 2450
<212> DNA
<213> Zea mays
```

400	21	ggcgggtcct	ccccggccaa	ggcgccggag	gggtccggcc	tcgaaccctc	ctcgctccgg	60
120	180	tcgcaggggc	gcaaggacca	gatacgccctc	ttcgacaagg	agttgctcca	ggttcggcac	120
180	240	caagttcata	tcggcgcgga	cgcggtccgc	aagaaaccgg	acatccctca	gaaagaaagg	180
240	300	atcaccccaa	tgctccaact	tgtgggtctt	gtctggccgg	agtaactcaa	gtcgggacct	240
300	360	gtctaacgca	ccctctggct	gcaggacagg	cccaaccgag	acatccacag	catccctgtac	300
360	420	gacgtgtttg	actaatctga	ggacgtccagg	gagcaggggc	gcccgggtgt	tgtgcatctgc	360
420	480	tcgcaggggg	tgtcacagctc	caagttccgtc	aaggcttgccc	gggggatctgc	aaatccaaac	420
480	540	caggagcttcg	atgacgacct	ccaagtctgc	cagaaagggtg	tgcatgctgat	tcggctgtca	540
540	600	atgggctttg	catgccagct	tcctccagtg	caggaaagggtg	tgcatgctgat	tcggctgtca	540
600	660	ccaaatccag	tgctcaggat	gtaacggcatg	ggcctccact	cccaagtatgc	ccctctgcat	600
660	720	ttgttgccca	aaatgctccta	tgacccatcc	ccaggccacc	ttgactctag	aggcgcgctc	660
720	780	atgttgcatg	ttctctcgtc	gctcctatgtc	tggtttggaa	tgaaagtgtga	tcgggttaatg	720
780	840	gaaaaggatg	caaggcgctg	tgctttccag	gtagttaggt	atgagaaagg	gcaggggcac	780
840	900	atcaaggttg	tgagagaaag	tcggagcg	caggagttct	gggatgcat	ttcaagtatg	840
900	960	ccaccataat	cagatagttaa	tacaagaatt	agcaaggacc	agatcgatc	agcatccaa	900
960	1020	agtgaaccag	gaagccggaa	aaatgagctc	tatgaagctg	attttgagct	tgtctacaaa	960
1020	1080	gcaatccactg	ggggagtgtg	ccctgcattt	tcaacctctg	gggtcgtgtga	tgagaccat	1020
1080	1140	cttccagcta	gagaaagttag	ctggaattta	ctgaggcaca	agtttatctc	caggttcgcta	1080
1140	1200	gctcgtgttt	atctcagatc	tgctcctaag	aaggattttg	atccacgggt	acaaacactg	1140
1200	1260	gctgcgtgag	catccaacctc	acctcccttc	cttctcccaa	gctccctatc	atcggatcca	1200
1260	1320	agtgtcagct	cgaagtatag	ttcagactca	ccctccctat	caacctacaac	tggtctctcca	1260
1320	1380	ccatcatctg	gctctcggcc	tgctctcatc	aatctggcac	atgcttttgt	ggcatcatcc	1320
1380	1440	aggtctcccc	ttctccaatc	atctaatgaa	ggagcttcaa	agcctctgg	catggaatca	1380
1440	1500	atacacctctc	cttcccaagac	ctctctcata	gcagaaaggaa	gaggagggct	cacactctca	1440
1500	1560	aagctaacctat	ctctccaaa	ggatcttgtta	ttggccaccaa	gggtggcgctc	tattgttatg	1500
1560	1620	ccaccaaagg	cggcatctcag	tattcggcagg	accggaggatg	cctcagatata	tagtatacaat	1560
1620	1680	gggtttaaaa	agctgaccta	tgagttttgc	tcagaaaaaat	gcactgtgtaa	tagttttgag	1620
1680	1740	tcgcattctg	aaactagatt	aattgagcgt	actgacagtta	actcagaaagt	ctgcagttaat	1680
1740	1800	gcacaacctg	tagtctacaa	gtggcccgcc	atgggaaagc	ttaactaat	tgcaaggcaag	1740
1800	1860	gatctctgaac	cgaagtctggt	tttaattttt	gttaacttcga	atggccatcag	gagaggagaa	1800
1860	1920	gcagttaaaa	tggtgtatat	atgggttagga	ggcgaaaaatg	agagcagcaca	gagtgttgac	1860
1920	1980	ttcgttcgatt	ggcaaacaggt	cactagtgtat	ttctctcatc	taaaaggcgct	cagcaaatgtt	1920
1980	2040	cttccctgtca	aggttttcaa	ggagcatgaa	gctgagaaatc	ttttggaaat	actgaaatgt	1980
2040		agtttaacatt	aggcagttagc	tatcaggata	atctgtagt	ctaaacaaa	tcgaaggagg	2040

catgccctcc agcatcagtc ggtaccgagtg atgtcagcg aggtataag ccacagccat 2100
tccctgaac ataataagct acaaacagat tccgttcgc aactgcgcct catgatctat 2160
atttgtcca gatggcagga ggcggccatg ggcttgtat cggttgcgaa ttagcactcg 2220
tggttagga gcaatcgcc gatcgggtg atatataccg ctccctgta atgtaagctc 2280
agatactgg agctggttg tcgacagta cttttagcc taacatctc tgcatactc 2340
tgaaaggac agattgtaa tcccttgac tatgtaatg gctccatgg tcataactc 2400
taaaccatgt ggaactcagt tgcctagctg taaaaaaaaa aaaaaaaaaa 2450

<210> 22
<211> 661
<212> PRT
<213> Zea mays

<400> 22
Gly Gly Ser Ser Pro Ala Lys Pro Gly Gln Gly Leu Arg Leu Asp Leu 1
5
10
15

Ser Ser Leu Arg Ser Gln Gly Arg Lys Asp Gln Ile Ala Phe Phe Asp 20
25
30

Lys Gln Cys Ser Lys Val Ala Asp His Val Tyr Leu Gly Gly Asp Ala 35
40
45

Val Ala Lys Asn Arg Asp Ile Leu Arg Lys Asn Gly Ile Thr His Val 50
55
60

Leu Asn Cys Val Gly Phe Val Cys Pro Gln Tyr Phe Lys Ser Asp Leu 65
70
75
80

Val Tyr Arg Thr Leu Trp Leu Gln Asp Ser Pro Thr Gln Asp Ile Thr 85
90
95

Ser Ile Leu Tyr Asp Val Phe Asp Tyr Phe Gln Asp Val Arg Gln Gln 100
105
110

Gly Gly Arg Val Leu Val His Cys Cys Gln Gly Val Ser Arg Ser Thr 115
120
125

Ser Leu Val Ile Ala Tyr Leu Met Trp Arg Gln Gly Gln Ser Phe Asp 130
135
140

Asp Ala Phe Gln Phe Val Lys Ala Ala Arg Gly Ile Ala Asn Pro Asn 145
150
155
160

Met Gly Phe Ala Cys Gln Leu Leu Gln Cys Gln Lys Arg Val His Ala 165
170
175

Ile Pro Leu Ser Pro Asn Ser Val Leu Arg Met Tyr Arg Met Ala Pro 180
185
190

His Ser Gln Tyr Ala Pro Leu His Leu Val Pro Lys Met Leu Asn Asp 195
200
205

Pro Ser Pro Ala Thr Leu Asp Ser Arg Gly Ala Phe Ile Val His Val 210
215
220

Leu Ser Ser Leu Tyr Val Trp Val Gly Met Lys Cys Asp Pro Val Met 225
 240
 Glu Lys Asp Ala Lys Ala Ala Phe Gln Val Val Arg Tyr Glu Lys 245
 250
 Val Gln Gly His Ile Lys Val Val Arg Glu Gly Leu Gln Pro Gln Glu 260
 265
 Phe Trp Asp Ala Phe Ser Ser Met Pro Pro Asn Ser Asp Ser Asn Thr 275
 280
 Lys Ile Ser Lys Asp Gln Ile Asp Ser Ala Ser Lys Ser Asp Pro Gly 290
 295
 Ser Arg Lys Asn Glu Ser Tyr Asp Ala Asp Phe Glu Leu Val Tyr Lys 305
 310
 Ala Ile Thr Gly Gly Val Val Pro Ala Phe Ser Thr Ser Gly Ala Gly 325
 330
 Asp Glu Thr His Leu Pro Ala Arg Glu Ser Ser Trp Ser Leu Leu Arg 340
 345
 His Lys Phe Ile Ser Arg Ser Leu Ala Arg Val Tyr Ser Asp Ser Ala 355
 360
 Leu Met Lys Asp Phe Asp Pro Arg Val Gln His Leu Ala Glu Ala 370
 375
 Ser Thr Ser Pro Pro Phe Leu Ser Pro Ser Ser Leu Ser Ser Asp Ser 385
 390
 Ser Val Ser Ser Lys Tyr Ser Ser Asp Ser Pro Ser Leu Ser Pro Thr 405
 410
 Thr Gly Ser Pro Pro Ser Phe Gly Leu Ser Pro Ala Ser Ser Asn Leu 420
 425
 Thr His Ala Leu Val Pro Ser Ser Arg Ser Pro Leu Ser Gln Ser Ser 435
 440
 Asn Glu Gly Ala Ser Lys Pro Ser Gly Met Glu Ser Ile His Ser Pro 450
 455
 Ser Lys Thr Ser Ser Ile Ala Glu Arg Arg Gly Gly Phe Thr Leu Leu 465
 470
 Lys Leu Pro Ser Leu Gln Lys Asp Leu Val Leu Pro Pro Arg Val Pro 485
 490
 Ser Ile Val Leu Pro Pro Arg Ala Pro Ser Ser Ile Arg Arg Thr Glu 500
 505
 Ser Ile Val Leu Pro Pro Arg Ala Pro Ser Ser Ile Arg Arg Thr Glu 510

Asn Cys Val Gly Phe Val Cys Pro Glu Tyr Phe Lys Ser Asp Phe Val
1
5
10
15

<400> 24
<213> Lycopersicon esculentum
<212> PRT
<211> 174
<210> 24

aactgtgtg ggttgtatg ccagagatg attcgtata ccgaacttg
120 tggtagag atagccatc agaaagatat actagatc tctatgagt ttttgactac
180 tttagagatg tcaggagca acatgggaag gttttgttc atgctgcca aggggtctc
240 cgtcaacct cgttgttat tgcctatcgt atgtggagag aaggacaaag ttttgatgat
300 gcccttgagt atgtaaagc agcaagggt atgctgcatc caaatatggg ttttgctgt
360 cagtattac aatgccaaa aagggttcat gcttctctc tgaagccaaag ttcattatta
420 agtagtaca gagtgcacc tcatccaaca tagatcctc tgcattcgt cccaaaatg
480 ttaaatgatc cctcaccgc agcatatgat tctagagtg catttatat acacataact
522 tcatcgtgat atgtatgat tggtaagaaa tgrgaagcaa tc

<400> 23
<213> Lycopersicon esculentum
<212> DNA
<211> 522
<210> 23

Leu Leu Asn Val Ser
660
Leu Pro Val Lys Val Phe Lys Glu His Glu Ala Glu Asn Leu Leu Glu
645 650 655
Gln Gln Val Thr Ser Asp Phe Leu His Leu Lys Gly Leu Ser Asn Val
625 630 635 640
Val Gly Gly Glu Asn Glu Ser Ser Lys Ser Val Asp Ser Val Asp Trp
610 615 620
Ser Asn Ala Ile Arg Arg Gly Glu Ala Val Lys Met Val Tyr Val Trp
595 600 605
Phe Ala Arg Lys Asp Leu Asp Pro Lys Ser Val Leu Ile Phe Val Thr
580 585 590
Ala Gln Leu Val Val Tyr Gln Trp Pro Ser Met Gly Lys Leu Thr
565 570 575
Thr Arg Leu Ile Glu Arg Thr Asp Ser Asn Ser Glu Val Cys Ser Asn
545 550 555 560
Phe Cys Ser Glu Lys Cys Thr Gly Asn Ser Leu Ser Ser His Ser Glu
530 535 540
Asp Ala Ser Asp Asn Ser Thr Asn Gly Val Lys Glu Leu Thr Ser Glu
515 520 525

<400> 26
ytcnkngcn ggnarrtngn tyt

<400> 25
gcnngcyltng crtytlytc c

Tyr Arg Thr Leu Trp Leu Gln Asp Ser Pro Ser Gln Asp Ile Thr Ser
20 25 30
Ile Leu Tyr Asp Val Phe Asp Tyr Phe Gln Asp Val Arg Gln His
35 40 45
Gly Lys Val Phe Val His Cys Cys Gln Gly Val Ser Arg Ser Thr Ser
50 55 60
Leu Val Ile Ala Tyr Arg Met Trp Arg Gln Gly Gln Ser Phe Asp Asp
65 70 75 80
Ala Phe Gln Tyr Val Lys Ala Ala Arg Gly Ile Ala Asp Pro Asn Met
85 90 95
Gly Phe Ala Cys Gln Leu Leu Gln Cys Gln Lys Arg Val His Ala Ser
100 105 110
Pro Leu Ser Pro Ser Ser Leu Leu Arg Met Tyr Arg Val Ala Pro His
115 120 125
Ser Pro Tyr Asp Pro Leu His Leu Val Pro Lys Met Leu Asn Asp Pro
130 135 140
Ser Pro Ala Ala Leu Asp Ser Arg Gly Ala Phe Ile Ile His Ile Pro
145 150 155 160
Ser Ser Val Tyr Val Trp Ile Gly Lys Lys Cys Gln Ala Ile
165 170

<220>
<223> Description of Artificial Sequence: degenerate
primer

<210> 26
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: degenerate
primer

<210> 25
<211> 21
<212> DNA
<213> Artificial Sequence

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 99/05413

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N9/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *

Citation of document, where appropriate, of the relevant passages

Relevant to claim No.

X

WO 94 23039 A (CANCER RES INST ROYAL
; MARSHALL CHRISTOPHER JOHN (GB); ASHWORTH
AL) 13 October 1994 (1994-10-13)
figure 7D
& DATABASE DGENE, online!

X

Accession no. R63602,
9 May 1995 (1995-05-09)
abstract

X

DATABASE SWISSPROT, online!
Accession no. P28562,
1 December 1992 (1992-12-01)
KEYES S.M. AND EMSLIE E.A.: "DUAL
SPECIFICITY PROTEIN PHOSPHATASE 1"
XP002131991
abstract

-/-

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not
considered to be of particular relevance

"E" earlier document but published on or after the international
filing date

"L" document which may throw doubts on priority claim(s) or
which is cited to establish the publication date of another
citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or
other means

"P" document published prior to the international filing date but
later than the priority date claimed

"T" later document published after the international filing date
or priority date and not in conflict with the application but
intended to understand the principle or theory underlying the
invention
"X" document of particular relevance; the claimed invention
cannot be considered novel or cannot be considered to
involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention
cannot be considered to involve an inventive step when the
document is combined with one or more other such docu-
ments, such combination being obvious to a person skilled
in the art.
"Z" document member of the same patent family

Date of the actual completion of the international search

1 March 2000

Date of mailing of the international search report

14/03/2000

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patendlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 851 epo nl,
Fax (+31-70) 340-3016

Authorized officer
Chakravarty, A

INTERNATIONAL SEARCH REPORT

Int. Application No	PCT/EP 99/05413
---------------------	-----------------

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
------------	---	-----------------------

X	DATABASE SWISSPROT 'online' Accession no. P28563, 1 December 1992 (1992-12-01) CHARLES C.H. ET AL.: "DUAL SPECIFICITY PROTEIN PHOSPHATASE 1 (EC 3.1.3.48) (EC 3.1.3.16)" XP002131992 cited in the application abstract	
P, X	DATABASE EMBL 'online' EBI accession no. AL085836, 28 June 1999 (1999-06-28) SALANUBAT M. ET AL.: "17 end of BAC F1115 of IGF library from strain Columbia of Arabidopsis thaliana" XP002131993 abstract	I-12
		I-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

Initial Application No
PCT/EP 99/05413

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
---	---------------------	----------------------------	---------------------

WO 9423039	A	13-10-1994	AU 677834 B	08-05-1997
			AU 6382394 A	24-10-1994
			CA 2157774 A	13-10-1994
			EP 0703984 A	03-04-1996
			JP 9501302 T	10-02-1997
			US 5958721 A	28-09-1999
			AU 696939 B	24-09-1998
			AU 1586195 A	29-08-1995
			CA 2182967 A	17-08-1995
			EP 0742827 A	20-11-1996
			WO 9521923 A	17-08-1995
			JP 9508795 T	09-09-1997

